



Data and Safety Monitoring Boards: Interim Monitoring for Clinical Trials

August 1, 2002

Sheryl F. Kelsey, Ph.D.

Department of Epidemiology

Graduate School of Public Health

University of Pittsburgh

DSMB is responsible for monitoring the trial

Data and Safety Monitoring Board (DSMB)

Primary charge: Oversee patient (participant) safety

particular issue in masked trials but even for unmasked trials investigators not informed as to overall results

DSMB - Description

- **Generally 3 – 10 members**
- **Members reflect disciplines needed to evaluate the trial**
 - **clinical**
 - **biostatistics**
 - **ethics**
 - **patient advocate (more recent/disease)**
- **No member associated with trial in any other capacity**
- **Appointed by sponsors of trial**
- **Ex-officio members**
 - NIH (if applicable)**
 - Industry sponsor (yes or no?)**
 - Trial Statistician**
 - Principal Investigator(s) (yes or no, masked?)**

DSMB's Responsibilities

Address ethical issues

Review protocol before initiation of trial

Review interim reports

recruitment

compliance

adverse effects

baseline comparability

treatment comparisons

Meet often enough to carry out responsibilities – in general once per year


additional meetings or conference calls if needed

Each meeting make recommendation that trial should stop, continue or continue w/modifications

Recruitment Reports

- Is the study or are the sites recruiting as many subjects as expected?
- If not, may recommend....
 - Remove sites
 - Add sites
 - Extend recruitment
 - Terminate trial

Cumulative Recruitment by Month As of September 28, 1998



Coordinating Center, University of Pittsburgh, Epidemiology Data Coordinating Center,
127 Parran Hall, 130 DeSoto Street, Pittsburgh, PA 15261, Telephone: 412/624-5447, FAX: 412/624-3775

- Confidential -
DSMB

Compliance

Assessment of compliance of...

treatments

inclusion of ineligible subjects

dropouts

completion of appropriate data collection
forms

certified assessors completing
assessments

proper treatment assignments

Adverse Effects

- **Dependent on type of study**
 - death
 - additional procedures
 - additional episodes or events
 - hospitalization
 - side effects (e.g. depression, nervousness, muscle aches)
- **By site (if multi-site)**
- **By treatment group**
- **For drug trials very specific FDA regulations**

Baseline Comparisons

Compare baseline characteristics of
population by treatment assignment
to: profile patient population
check balance

Treatment Comparisons

- Compare the treatments the trial to see if there are any differences
- Problem: If each comparison has a Type I error rate of .05, the chances of finding a false significant difference increase (classical statistical theory)

Example

2.5 years recruitment

One year follow-up


Treatment comparisons will be conducted every six months

- 1st analysis: 20 % of subjects complete year 1 visit
- 2nd analysis: 40% of subjects complete year 1 visit
- 3rd analysis: 60% of subjects complete year 1 visit
- 4th analysis: 80% of subjects complete year 1 visit
- 5th analysis: 100% of subjects complete year 1 visit

Example

- Total of 5 analyses
- If the Type I error rate at each of the analyses is .05, then the true Type I error rate is...

Type I Error Rates



- **Several methods for adjusting the Type I error rate**
- **Using these methods the Type I error rate is adjusted at each of the interim analyses so an overall Type I error rate of .05 can be maintained.**

Pocock

- **Reference:**
Pocock SJ. Group sequential methods in the design and analysis of clinical trials. Biometrika, 1977; 64:191-199.
- **Basic concept: keep the adjusted Type I error rate at each of the interim analyses the same (but less than .05) to maintain the overall Type I error rate of .05.**


Pocock

	Number of Looks			
	2	3	4	5
Adjusted Type I error rate	.0294	.0221	.0182	.0152
Z	2.178	2.281	2.361	2.413

For a normally distributed outcome. Where
Z=Value from standard normal distribution
corresponding to adjusted Type I error rate

Pocock

Significance Bounds



For 5 analyses

Pocock

- Problem: With constant adjusted Type I error rate, we don't have much Type I error left for the final comparison.
- Solution: “Spend” little of the Type I error at the earlier analyses, saving the majority for the final analysis.

O'Brien & Fleming

- **Reference:**
O'Brien PC and Fleming TR. A Multiple Testing Procedure for Clinical Trials. Biometrics, 1979; 35:549-556.
- **Basic concept:** Keep the adjusted Type I error rate small early on, increasing the adjusted Type I error rate over time, maintaining the overall Type I error rate.

O'Brien & Fleming example

	Look Number				
	1	2	3	4	5
Adjusted Type I error rate	.00128	.00144	.00164	.00219	.04806
Z	3.00	2.98	2.94	2.85	1.67


Where:

5 planned looks

Z=Value from Standard Normal distribution
corresponding to adjusted Type I error rate

O'Brien & Fleming

Significance Bounds



For 5 analyses (looks)

O'Brien & Fleming

- Problem: What if recruitment is slower than expected, or the Safety and Data Monitoring Board Requests more analyses. The two methods presented thus far are designed for planned interim analyses.
- Solution: A spending function that does not depend on the total number of analyses.

Lan & DeMets

- Reference

Lan KKG and DeMets DL. Discrete sequential boundaries for clinical trials. Biometrika, 1983;70:659-663.

- Basic concept: The Type 1 error is spent on a function that is dependent on past and current analysis times, not future times or the total number of analyses.
- “Spending the alpha”

Other Monitoring Tools

Bayesian Methods

- Assessment of evidence does not depend on plan for future analysis (no “alpha spending”)

Conditional Power

- What is the probability that a difference will be detected at the end given the data so far.

DSMB Decision Making Process

- Decision to stop a trial not purely statistical
- Magnitude of difference must be considered
- Possible subgroup effects
- Should the protocol be modified?

Some Reasons Why a Clinical Trial Might be Stopped Early

- Treatments are found to be convincingly different by impartial knowledgeable experts.
- Treatments are found to be convincingly not different by impartial knowledgeable experts.
- Side effects or toxicity are too severe to continue treatment in light of potential benefits.
- The data are poor quality
- Accrual is too slow to complete the study in a timely fashion.
- Definitive information becomes available from outside the study, making the trial unnecessary or unethical.
- The scientific question are no longer important because of other developments.
- Adherence to the treatment is unacceptably poor, preventing an answer to the basic question.
- Resources to perform the study are lost or no longer available.
- The study integrity has been undermined by fraud or misconduct.

Disadvantages of Stopping Early

- **Lack of credibility**
small trials not convincing
- **Bias/Mistake**
Trial liable to stop on “random high”, risk of false positive
- **Pressure**
unduly enthusiastic and extrapolated recommendation may follow
- **Long term outcome not well known**

Confidentiality of Interim Reports

May change investigators outlook and participation based on results that are not complete

- Should industry sponsors/supporters have access?**

Turn in reports at the end of the meeting

- **Though DSMB members are very interested in the science and can make valuable suggestions for the design execution and analysis of a trial, must keep in mind that patient safety is primary focus to ensure an ethical trial**